


# Diabetes Prevalence\* in First Nations Communities by NIHB region





- Diabetes prevalence was not very different between the four NIHB regions
- However, all regions had higher prevalence than the overall Ontario population


\*age- and sex-standardized rates for people residing in First Nations Community RESCODES, March 31, 2012

\*\*Ontario rates are for 2011

# Diabetes Incidence\* in First Nations Communities by NIHB region



- Moose Factory Region
- ▲ Southern Ontario Region
- Sioux Lookout Region
- ◆ Thunder Bay Region






- Diabetes incidence was higher in all NIHB regions compared to the overall Ontario population

\*age- and sex-standardized rates for people residing in First Nations Community RESCODES, April 1, 2011 - March 31, 2012

\*\*Ontario rates are for April 1, 2009 - March 31, 2010

# Diabetes Skin and Soft Tissue Complications\* in First Nations Communities by NIHB Region



- Rates of hospital or emergency department visits for diabetes-related skin and soft tissue infection or foot ulcer were higher in all NIHB regions compared to the overall Ontario population

- Sioux Lookout Region
- Moose Factory Region
- ▲ Southern Ontario Region
- ◆ Thunder Bay Region

\*age- and sex-standardized rates for people residing in First Nations Community RESCODES, April 1, 2007 - March 31, 2010  
 \*\*Ontario rates are for April 1, 2006 - March 31, 2011